

DOI: 10.19177/978-85-8019-206-3.23-41

Foraminifera biodiversity coupled with environmental quality in Pium River estuary and Pirangi Coral Reef (RN, Brazil)

A biodiversidade foraminifera associada à qualidade ambiental no estuário do Rio Pium e no Coral de Pirangi (RN, Brasil)

Patrícia Pinheiro Beck Eichler¹

Cristiane Leão Cordeiro de Farias²

Diogo Santos³

Audrey Amorim⁴

Alison Mendonça⁵

Helenice Vital⁶

Moab Praxedes Gomes⁷

ABSTRACT: The quality assessment of coastal and estuarine zone of Pium River Estuary and Pirangi coral reef area (inner shelf of Rio Grande do Norte) in recent and past environments is studied by organisms capable of: synthesizing the environmental characteristics (oceanographic and geological), highlighting the environmental variations of short periods and reacting sensitively to seasonal variations and anthropogenic effects. This work relates the ecological indices of the foraminifers' indicators species and the dominance of *Ammonia tepida*, *Bolivina striatula*, *Discorbis peruvianus*, *Elphidium articulatum*, *E. discoideale*, *Quinqueloculina lamarckiana*, *Q. intricata*, *Q. patagonica*, *Textularia gramen* and *T. earlandi* to characterize ecologically different environments as means of salinity, temperature and sediment characteristics, as well as levels of industrial pollution and organic contamination in the last

1 Bióloga, Doutora em Oceanografia Biológica, Programa de Pós-graduação em Ciência Ambiental, Univ. do Sul de Santa Catarina (PPGCA/UNISUL), Programa de Pós-graduação em Geofísica e Geodinâmica Univ. Federal do Rio Grande do Norte

2 Bióloga, Mestre em Geodinâmica, Univ. Federal do Rio Grande do Norte (PPGG/UFRN), crislc83@yahoo.com.br

3 Geólogo, Mestre em Geodinâmica, Univ. Federal do Rio Grande do Norte (PPGG/UFRN), diogo_geologia@hotmail.com

4 Bióloga, Bacharel, Programa de Pós-graduação em Ciência Ambiental, Universidade do Sul de Santa Catarina (PPGCA/UNISUL), audreyamorim-correa@gmail.com

5 Geóloga, Doutora em Geologia Marinha, Programa de Pós-graduação em Geofísica e Geodinâmica Univ. Federal do Rio Grande do Norte (PPGG/UFRN), helenicevital2004@yahoo.com.br

6 Geólogo, Doutora em Geologia Marinha, Programa de Pós-graduação em Geofísica e Geodinâmica, Univ. Federal do Rio Grande do Norte (PPGG/UFRN), gomesmp@geologia.ufrn.br

7 asdfasdfsdfasfd asdfasdfsdf asdfasdfsdf asdfasdfsdf

25 years. The three shallow sediment cores analyzed have showed higher species diversity and evenness about 18-20 years ago followed by recent periods with greater dominance and environmental instability. We observed inter annual fluctuations but there is a trend of decreasing foraminifera diversity and evenness, with greater dominance of less foraminifera species nowadays very close to fragile environments, the coral reef area and the estuary of the Pium River in a carbonate shelf. Our results indicate a decrease in biodiversity in the whole ecosystem towards today.

■ **Key words:** Diversity. Evenness. Dominance. Assessment. Coastal. Industrial pollution. Organic contamination.

RESUMO: A avaliação da qualidade da zona costeira e estuarina do Estuário do Rio Pium e da área de recifes de coral de Pirangi (plataforma interna do Rio Grande do Norte) em ambientes recentes e passados é estudada por organismos capazes de sintetizar as características ambientais (oceanográficas e geológicas), destacando variações ambientais de curtos períodos e reagindo sensivelmente a variações sazonais e efeitos antrópicos. Este trabalho relaciona os índices ecológicos de espécies indicadoras de foraminíferos e a dominância de *Ammonia tepida*, *Bolivina striatula*, *Discorbis peruvianus*, *Elphidium articulatum*, *E. discoidale*, *Quinqueloculina lamarckiana*, *Q. intricata*, *Q. patagonica*, *Textularia gramen* e *T. earlandi* para caracterizar ambientes ecologicamente diferentes como respeito à salinidade, temperatura e características de sedimentos, bem como níveis de poluição industrial e contaminação orgânica nos últimos 25 anos. Os três sedimentos superficiais analisados apresentaram maior diversidade de espécies e uniformidade há 18-20 anos, seguidos por períodos recentes com maior dominância e instabilidade ambiental. Observamos flutuações interanuais, mas há uma tendência de diminuir a diversidade e a estabilidade das espécies de foraminíferos, com aumento de dominância, em direção aos dias atuais em ambiente muito frágil, os recifes coralíneos, o estuário do Rio Pium, em uma plataforma carbonática. Nossos resultados indicam uma diminuição na biodiversidade em todo o ecossistema em relação a hoje.

■ **Palavras-chave:** Diversidade. Estabilidade. Dominância. Avaliação. Costeira. Poluição industrial. Contaminação orgânica.

1 INTRODUCTION

The use of biological indicators fossils (foraminifera), a simple handling and low cost tool is a good alternative for environmental studies in biodiversity, oceanography and environmental monitoring, as they are plentiful and easy to collect, present in all aquatic environments. Assessment and diagnosis of the carbonate inner shelf of Pium River Estuary and adjacent reef area in Pirangi (Rio Grande do Norte) in recent and past environment is being accomplished through these organisms capable of synthesizing general characteristics of the environment (oceanographic and geological), highlighting the environmental changes in short periods and react sensitively to seasonal

variations and anthropogenic effects. The works that deal with the distribution of these fossils indicators aim to characterize different environments as salinity, temperature and sediment characteristics, as well as levels of industrial pollution, organic pollution⁸. Their potential for fossilization provides data to understand the current environmental quality of the region. Our objectives are to evaluate if there is a shift of biodiversity due to anthropogenic stress, and to verify if there are interannual changes in the past 25 years.

2 METHODOLOGY

Three (3) shallow cores (about 25cm) were collected along the coral reef southward. Core 1 was closest to the river Pium and the other two were collected at 500m away, as shown in Figure 1.

Figure 1 – Location and photograph of open cores (T1, T2 and T3) collected along the reef area.



- 8 EICHLER, P.P.B.; EICHLER, B.B.; MIRANDA, L.B. de; RODRIGUES, A.R. Foraminiferal assemblages in a subtropical, mixohaline, estuarine channel, Bertioiga (São Paulo). *Braz. J. Res.*, n. 37, p. 45-58, 2007; EICHLER, P.P.B.; SEN GUPTA, B.K.; EICHLER, B.B.; BRAGA, E.S.; CAMPOS, E.J. Benthic foraminiferal assemblages of the South Brazil: Relationship to water masses and nutrient distributions. *Cont. Shelf. Res.*, n. 28, p. 1674-1686, 2008; EICHLER, P.P.B.; BILLUPS, K.; VELEZ, C.C. Investigating faunal and geochemical methods for tracing salinity in an Atlantic Coastal Lagoon, Delaware, USA. *J. For. Res.*, n. 40, p. 14-33, 2010; EICHLER, P.P.B.; EICHLER, B.B.; SEN GUPTA, B.K.; RODRIGUES, A.R. Foraminifera as indicators of marine pollutant contamination on the inner continental shelf of southern Brazil. *Marine Pollution Bulletin*, n. 64, p. 22-30, 2012; YANKO, V.; KRONFELD, J.; FLEXER, A. Response of benthic foraminifera to various pollution sources: implications for pollution monitoring. *J. For. Res.*, n. 24, p. 1-17, 1994; YANKO, V. Benthic foraminifera as bioindicators of stress environment: anthropogenic problems – foraminiferal solution. In: *The First International Conference Application of Micropaleontology in Environmental Sciences*. Tel Aviv, Herzlia: Anamet, 1997. p.117-119; YANKO, V.; AHMAD, M.; KAMINSKY, M. Morphological deformities of benthic foraminiferal tests in response to pollution by heavy metals: implications for pollution monitoring. *J. For. Res.*, n.28, p. 177-200, 1998; YANKO, V.; ARNOLD, A.J.; PARKER, W.C. Effects of marine pollution on benthic Foraminifera In: SEN GUPTA, B.K. (Ed.). *Modern Foraminifera*. New York: Kluwer Acad. Publ., 1999. p. 217-235.

The study area is located in the coastal zone of the Estuary Pium and inner shelf adjacent to the reef area Pirangi. The cores were collected in April 2012 in a depth of 6 meters. After drying for a few weeks they were cut in half and photographed. Then sub sampling was performed for the fauna of foraminifera every two centimeters. The cores had recovered the following sizes: 20 cm with T1, T2 and T3 with 24 cm each. Based on the rate of sedimentation in coastal internal platforms⁹ we can approach that 1 cm is equals to about one year in coastal cores.

Each sample was washed and sieved into two sieves of 0.500 mm and 0.062 mm. The material retained on the sieves was dried on filter paper and then placed in an oven at 50 degrees for 72 hours. The material of the 0.062 mm sieve went through the process of splitting for later sorting, identification, counted and classification of species of benthic foraminifera in stereoscope. During the separation of each sample, the specimens are transferred with the aid of special brush to slides with black background. Absolute frequency tables of foraminiferal species were made. Univariate (diversity indices, evenness and dominance) and multivariate (Cluster and MDS) analysis were applied using Primer 6 program.

3 RESULTS

Core 1 shows the presence of 57 species with mainly occurrence of *Ammonia tepida*, *Discorbis peruvianus*, *Elphidium articulatum*, *Quinqueloculina lamarckiana*, *Q. intricata*, *Q. patagonica*, and *Textularia gramen* in all times (Table 1).

Table 1 – Absolute frequency of species of foraminifera identified in Core 1.

Species/Stations	1 (top)	2	3	4	5	6	7	8	9	10 (bottom)
<i>Ammonia rolshauseni</i>	0	0	200	180	45	0	0	560	100	420
<i>Ammonia tepida</i>	4130	4725	920	1215	405	3255	1320	5800	380	7080
<i>Amphisorus sp.</i>	0	0	0	0	0	0	0	0	0	30
<i>Amphistegina sp.</i>	0	0	0	0	0	0	0	0	20	30
<i>Arenoparrella mexicana</i>	35	175	40	90	0	0	0	0	0	180

9 PATCHINEELAM, S.; SMOAK, J. Sediment Accumulation Rates along the Inner Eastern Brazilian Continental Shelf. *Geo-Marine Letters*, n. 19, p. 196, 1999. Disponível em: <<https://doi.org/10.1007/s003670050109>>.

Species/Stations	1 (top)	2	3	4	5	6	7	8	9	10 (bottom)
<i>Bolivina difformis</i>	210	140	0	45	0	0	0	80	0	180
<i>Bolivina striatula</i>	700	665	0	0	90	560	0	400	80	1080
<i>Bolivina sp.</i>	0	0	0	45	0	0	0	80	0	0
<i>Bolivina tortuosa</i>	0	175	0	135	0	0	0	40	20	150
<i>Bolivina translucens</i>	245	210	0	0	45	0	120	0	0	0
<i>Bulimina patagonica</i>	0	0	0	0	0	35	0	40	20	120
<i>Bulimina sp.</i>	0	0	0	0	0	0	0	0	0	30
<i>Buccella peruviana</i>	0	385	320	45	315	140	160	360	120	480
<i>Bulimina marginata</i>	0	70	0	0	0	0	0	0	0	30
<i>Buliminella elegantissima</i>	140	140	0	0	0	0	0	0	0	120
<i>Cassidulina subglobosa</i>	140	70	0	0	0	0	0	80	0	60
<i>Cibicides dispars</i>	70	0	0	0	45	0	0	0	0	0
<i>Cibicides fletcheri</i>	35	0	0	0	90	0	0	80	0	120
<i>Cibicides variabilis</i>	0	70	0	0	45	0	0	0	0	30
<i>Cornuspira involvens</i>	665	210	0	45	45	175	40	40	20	450
<i>Discorbis peruvianus</i>	560	630	80	315	225	385	80	840	100	1620
<i>Discorbis valvulatus</i>	0	0	200	135	0	0	40	200	240	120
<i>Discorbis williamsoni</i>	70	175	600	135	0	0	80	560	0	300
<i>Elphidium articulatum</i>	70	175	40	225	225	210	200	240	140	60
<i>Elphidium discoidale</i>	280	105	0	0	135	315	160	120	0	180
<i>Elphidium galvestonense</i>	0	0	120	90	0	0	0	200	20	150
<i>Elphidium magellanicum</i>	70	0	0	0	0	0	0	120	20	90
<i>Elphidium sp.</i>	35	35	0	0	0	0	0	0	0	30
<i>Fissurina laevigata</i>	35	0	0	0	0	0	0	0	0	30
<i>Fissurina lucida</i>	70	70	0	0	0	0	0	0	0	30
<i>Hanzawaia boueana</i>	70	70	80	90	45	35	80	160	0	270
<i>Hopkinsina pacifica</i>	0	105	0	45	0	35	40	40	0	0
<i>Miliolinella subrotunda</i>	0	0	1000	1125	135	0	40	840	160	161

Species/Stations	1 (top)	2	3	4	5	6	7	8	9	10 (bottom)
<i>Orthomorphina sp.</i>	0	0	40	0	0	35	40	160	60	120
<i>Patellina corrugata</i>	0	0	0	0	0	0	0	0	0	60
<i>Poreponides lateralis</i>	280	210	80	90	0	35	200	120	40	360
<i>Pseudononium atlanticum</i>	105	70	0	90	45	70	0	80	40	240
<i>Pyrgo nasuta</i>	35	35	80	45	0	35	0	40	0	30
<i>Pyrgo ringens</i>	0	70	400	270	270	70	40	120	20	90
<i>Pyrgo subsphaerica</i>	0	0	0	0	45	0	40	0	0	0
<i>Quinqueloculina gregaria</i>	35	35	0	135	0	35	0	200	0	0
<i>Quinqueloculina intricata</i>	105	280	280	225	225	35	80	360	40	420
<i>Quinqueloculina lamarkiana</i>	2835	2730	3160	2295	990	1505	1000	4360	620	3930
<i>Quinqueloculina patagonica</i>	2170	2240	560	1305	1620	1715	960	1560	160	1650
<i>Quinqueloculina seminulum</i>	0	0	0	270	0	0	0	0	40	120
<i>Quinqueloculina sp.</i>	70	0	40	0	360	35	40	0	20	120
<i>Textularia earlandi</i>	0	0	80	45	90	70	120	80	40	120
<i>Textularia gramen</i>	35	525	560	585	675	420	400	840	240	720
<i>Triloculina baldai</i>	70	420	0	0	135	70	40	80	20	30
<i>Trochammina inflata</i>	0	35	120	0	0	0	0	80	0	0
<i>Trochammina ochracea</i>	70	35	40	0	0	0	0	0	0	30
<i>Trochammina plana discorbis</i>	70	35	80	45	0	0	120	160	40	0
<i>Trochammina sp.</i>	0	0	0	45	0	0	0	0	0	30
<i>Trochammina squamata</i>	35	35	0	45	0	35	0	160	0	60
<i>Uvigerina peregrina</i>	35	35	0	0	0	0	0	0	0	60
<i>Uvigerina striata</i>	0	70	0	45	0	0	0	120	0	330
<i>Wiesnerella sp.</i>	0	35	0	45	0	210	0	0	0	90


Source: authors

It is observed that the stations 9 and 10 show higher diversity and evenness followed by stations 4 and 5. Interval 6 and 1 (top) show the highest values of dominance and lower diversity and evenness.

Table 2 – Intervals, Number of species (S), number of specimens (N), Evenness, Diversity, and Dominance in Core 1

Intervals	(S)	(N)	Evenness	Diversity	Dominance
1 (top)	33	13580	0.65	2.29	0.17
2	39	15365	0.67	2.47	0.15
3	25	9520	0.76	2.44	0.14
4	34	9720	0.76	2.68	0.11
5	25	6840	0.81	2.63	0.10
6	26	9765	0.68	2.23	0.17
7	25	5760	0.77	2.50	0.12
8	39	19920	0.69	2.55	0.14
9	28	3060	0.83	2.79	0.08
10 (base)	52	22931	0,67	2,68	0,14

Source: authors


Figure 2 – Evenness, Diversity and Dominance in Core 1

Source: authors

The groups formed by the cluster analysis and the definition of MDS showed 3 main groups (10, 1 & 2), (3, 4, and 8) (5, 6, 7, and 9) (Figure 3). Group 1 formed by the most recent to oldest stations (1 and 2) and the interval 10. This group may have been formed by high dominance of a few species such as: *Ammonia tepida*, *Cornuspira involvens*, and the appearance of rare species like *Buliminella elegantissima*, *Fissurina lucida*, *F. laevigata*, *Uvigerina peregrina*, and *Elphidium* sp. The other two groups formed are not

clear about the group. This core by being closer to the estuary of the River Pium may be showing an influence of seasonal freshwater discharge.

Figure 3 – Cluster analysis and MDS in Core 1



Source: authors

Analyzing Table 3 absolute frequency of core 2 shows the presence of 39 species and the occurrence of *Ammonia tepida*, *Bolivina striatula*, *Discorbis peruvianus*, *Elphidium discoideale*, *Quinqueloculina Lamarckian*, *Q. patagonica*, and *Textularia earlandi* in all periods since the most recent to the most current. Compared to core 1, it is observed that the core 2 shows a much smaller number of specimens. One can note a few species at the top (1) with a predominance of *Quinqueloculina lamarckiana*. (2) The interval features three dominant species: *Ammonia tepida*, *Quinqueloculina lamarckiana* and *Q. patagonica*. In Sample 3 besides these three species the interval 2 also present *Bolivina striatula* and *Discorbis peruvianus* with relatively high frequency. In interval 4, *Discorbis williamsoni* also appears in representative number. Intervals 5 and 6 show a predominance of *A. tepida*. Sample 7 shows the same in the other abundant species, and contain *Textularia earlandi* and *Milionella subrotunda* in significant numbers. Highlighting the large number of specimens found in this season. Samples 9 and 10 retain the same characteristics of the other, with no major highlights. Already the 11 sample stands out as the most diverse when compared with others, as well as the large amount of specimens. The base (12) shows a small number of specimens of *A. tepida*.

Table 3 – Absolute frequency of species of foraminifera identified in Core 2

Species/Stations	1 (top)	2	3	4	5	6	7	8	9	10	11 (bottom)	12
<i>Ammonia rolshauseni</i>	0	0	1	0	0	0	0	0	1	0	0	2
<i>Ammonia tepida</i>	8	23	44	57	88	87	111	74	37	57	121	13
<i>Amphicoryna scalaris</i>	0	0	1	2	1	2	0	0	2	2	0	0
<i>Bolivina striatula</i>	2	4	11	18	15	28	28	30	15	24	60	28
<i>Bolivina tortuosa</i>	0	0	0	0	5	5	2	1	0	0	1	1
<i>Buccella peruviana</i>	0	2	0	0	1	6	4	6	9	20	24	26
<i>Bulimina marginata</i>	0	1	3	0	0	0	0	0	0	1	3	0
<i>Cassidulina subglobosa</i>	1	0	0	2	0	3	5	5	1	1	4	1
<i>Cibicides fletcheri</i>	0	3	1	0	1	0	0	0	0	0	3	0
<i>Cibicides variabilis</i>	2	0	0	0	1	1	1	0	0	2	1	3
<i>Cornuspira envolvens</i>	1	1	1	2	2	3	1	5	9	3	12	1
<i>Discorbis peruvianus</i>	1	1	4	15	13	28	33	17	11	15	29	22
<i>Discorbis valvulatus</i>	0	1	4	2	7	1	0	2	0	0	1	3

<i>Discorbis williamsoni</i>	1	1	1	10	0	1	2	5	0	1	1	0
<i>Elphidium discoidale</i>	1	3	1	4	6	1	4	4	0	1	1	3
<i>Elphidium poeyanum</i>	0	0	1	0	0	0	0	2	2	0	0	0
<i>Fissurina laevigata</i>	0	0	1	0	2	0	3	1	1	0	0	1
<i>Gaudryna exilis</i>	0	0	0	1	0	2	0	0	0	0	0	0
<i>Hanzawaia boueana</i>	1	2	0	4	0	2	4	6	0	0	1	0
<i>Hopkinsina pacifica</i>	0	0	0	0	0	0	1	2	0	0	1	0
<i>Lagena striata</i>	0	0	0	0	0	0	0	1	0	0	1	0
<i>Milliolina subrotunda</i>	0	0	2	0	1	0	7	2	2	1	2	7
<i>Patellina corrugata</i>	0	1	0	5	5	12	4	3	1	0	12	8
<i>Planorbilina mediteranensis</i>	0	0	0	0	0	1	1	0	0	0	0	0
<i>Poroeponides lateralis</i>	1	0	4	1	2	0	0	0	1	0	1	3
<i>Pseudonionium atlanticum</i>	0	0	0	4	2	2	3	1	0	1	3	5
<i>Pyrgo nasuta</i>	1	0	0	0	1	2	0	1	0	0	3	0
<i>Quinqueloculina angulata</i>	0	0	0	0	0	0	2	1	1	0	0	0
<i>Quinqueloculina intricata</i>	0	1	0	0	2	1	3	0	2	0	3	1
<i>Quinqueloculina lamarckiana</i>	34	21	54	43	48	38	44	32	17	12	63	69
<i>Quinqueloculina patagonica</i>	9	23	16	21	22	43	37	22	18	19	58	23
<i>Quinqueloculina sp</i>	0	3	0	0	1	3	2	2	2	2	2	3
<i>Robalus rotulatus</i>	1	0	3	0	0	0	0	0	0	0	0	0
<i>Spirilocolina planulata</i>	0	0	1	0	0	1	0	0	0	0	0	0
<i>Textularia earlandi</i>	3	0	3	8	5	7	10	4	9	2	13	6
<i>Trilocolina baldai</i>	0	0	4	1	2	9	1	1	0	0	0	0
<i>Trochammina ochracea</i>	0	1	1	3	1	3	1	5	1	0	4	2
<i>Trochammina squamata</i>	0	0	0	0	0	13	4	0	0	0	0	0
<i>Uvigerina peregrina</i>	1	0	0	1	0	2	0	0	0	1	1	0

Source: authors

Table 4 shows the number of species varied between 16 and 29 and the number of species varied between 68 and 430 individuals. Evenness showed a lower value (0.66) in the interval 3 and higher value (0.79145) in the interval 9. Diversity was lower in the interval 1 (1.83) and highest in the interval 6 (2.43). The dominance showed the lowest value (0.12) in the interval 9 and the highest value in the interval 1 (0.28)


Table 4 – Number of species (S), individuals (N), evenness, diversity, and dominance

Intervals	(S)	(N)	Evenness	Diversity	Dominance
1	16	68	0.66	1.83	0.28
2	17	92	0.72	2.06	0.18
3	22	12	0.67	2.08	0.20
4	20	24	0.75	2.20	0.15
5	24	24	0.66	2.12	0.20
6	28	37	0.72	2.43	0.13
7	26	38	0.68	2.24	0.17
8	26	25	0.73	2.38	0.15
9	20	12	0.79	2.37	0.12
10	18	15	0.71	2.06	0.18
11	29	430	0.68	2.29	0.14
12	22	231	0.76	2.36	0.13

Source: authors

From the analysis of Figure 4 can be noted the interval 1 (top) because of lower diversity and higher dominance. The interval 6 shows the greatest diversity of core. Stations 4 and 9 show the highest indices of evenness which shows no species dominating over another exists. One can also notice low levels of dominance.


Figure 4 – Data diversity, dominance and evenness of Core 2



Source: authors

According to the number of species (Figure 5), it can be seen that several stations were the number 6 and 11 opposed, leaving the top station as less diverse.


Figure 5 – Number of species of foraminifera Core 2



Source: authors

Figure 6 shows the number of individuals found at each station, revealing that the sample 11 has the largest number of subjects followed by the stations 6 and 7. Again, the sample is revealed as the top with a lower number of specimens.


Figure 6 – Number of individuals per station



Source: authors

From Figure 7a observed the formation of two major groups excluding the interval 5. From Figure 7b we find that the interval 1 (top) was less diverse, followed by samples 2, 3, 4, 5. Intervals 9, 10, 11 and 12 had the highest diversity.

Figure 7a – Cluster analysis among species of foraminifera and Figure 7b – Analysis of MDS stations.



We can observe the presence of 70 species and the occurrence of *Ammonia tepida*, *Elphidium discoidale*, *Quinqueloculina lamarckiana*, and *Q. patagonica*, in all periods.

Table 5 – Absolute frequency of foraminiferal species in core 3


Species/Stations	1 (top)	2	3	4	5	6	7	8	9	10	11	12 (bottom)
<i>Agglutinella agglutinans</i>	0	0	0	120	0	0	0	60	0	100	0	0
<i>Ammonia tepida</i>	5370	6630	15500	26160	80	668	33540	15000	495	11050	598	5940
<i>Arenoparrella mexicana</i>	0	210	0	200	0	0	0	0	0	150	0	0
<i>Bolivina sp.</i>	0	0	0	0	0	0	0	60	0	100	0	0
<i>Bolivina striatula</i>	240	270	1175	1000	6	69	0	1080	90	600	26	360
<i>Bolivina tortuosa</i>	0	270	200	1040	0	0	120	420	0	600	0	45
<i>Buccella peruviana</i>	0	0	0	40	0	0	0	240	990	600	17	0
<i>Bulimina marginata</i>	0	120	0	440	0	5	0	300	0	0	0	45
<i>Bulimina patagonica</i>	0	120	425	360	0	9	600	240	0	300	0	45
<i>Buliminella elegantissima</i>	0	180	0	0	0	0	0	120	0	50	0	0
<i>Cassidulina subglobosa</i>	0	0	0	40	0	0	0	600	0	250	0	45
<i>Cibicides fletcheri</i>	0	180	0	320	0	0	0	300	0	400	0	0
<i>Cibicides variabilis</i>	0	30	0	0	0	0	0	0	0	500	0	135
<i>Cornuloculina sp.</i>	0	0	0	0	0	0	0	60	0	50	0	0
<i>Cornuspira involvens</i>	60	60	200	160	1	0	120	120	0	550	13	90
<i>Discorbis peruvianus</i>	0	450	75	2120	7	76	540	1920	45	1900	8	315
<i>Discorbis valvulatus</i>	0	0	0	0	0	0	0	60	0	50	0	0
<i>Discorbis williansoni</i>	0	90	0	80	0	0	360	120	0	50	0	45
<i>Elphidium articulatum</i>	0	0	125	360	0	0	0	180	135	150	25	0
<i>Elphidium discoidale</i>	1110	120	900	400	17	125	1680	360	360	100	8	45
<i>Elphidium excavatum</i>	0	60	0	160	0	0	0	180	0	400	0	0
<i>Elphidium galvestonense</i>	0	0	0	680	0	0	0	60	0	100	0	0
<i>Elphidium sp.</i>	0	90	0	80	0	0	0	0	0	0	0	0
<i>Falsagglutinella angularis</i>	0	30	0	0	0	0	0	60	0	0	0	0
<i>Fissurina laevigata</i>	30	30	75	0	0	1	0	0	0	0	9	0
<i>Fissurina lucida</i>	0	0	0	40	0	0	120	0	0	0	0	0
<i>Floresina spicata</i>	30	0	0	0	0	0	0	0	0	50	0	0
<i>Fursenkoina sp.</i>	0	0	25	40	0	0	0	0	0	0	0	0
<i>Hanzawaia boueana</i>	0	0	0	560	0	0	0	300	0	250	0	180
<i>Hopkinsina pacifica</i>	0	30	0	0	0	0	0	180	0	100	0	0
<i>Lagena laevis</i>	0	0	0	0	0	0	0	0	150	1	0	0
<i>Lagena striata</i>	0	0	0	80	0	0	60	0	0	0	0	0
<i>Lobatula lobatula</i>	0	0	0	0	0	0	0	0	100	0	0	0
<i>Loxostomina sp.</i>	0	0	0	0	0	0	120	0	0	0	0	0
<i>Massilina planata</i>	0	0	0	0	0	0	60	0	100	0	0	0

Species/Stations	1 (top)	2	3	4	5	6	7	8	9	10	11	12 (bottom)
<i>Massilina timorensis</i>	0	0	0	40	0	0	0	0	0	0	0	0
<i>Miliolinella subrotunda</i>	90	540	25	1600	0	16	480	1680	0	2950	36	1350
<i>Patellina corrugata</i>	0	0	0	0	0	0	0	0	0	150	0	45
<i>Peneroplis jovem</i>	0	0	25	0	0	0	0	60	0	0	0	0
<i>Peneroplis pertusus</i>	0	0	0	240	0	0	0	180	0	200	0	0
<i>Poroeponides lateralis</i>	0	90	25	160	0	0	0	60	0	50	0	0
<i>Pseudonionium atlanticum</i>	30	30	175	80	0	3	240	360	0	550	33	45
<i>Pseudotriloculina patagonica</i>	0	0	0	0	0	0	0	0	0	150	0	405
<i>Pyrgo nasuta</i>	30	0	0	0	0	0	0	60	0	0	0	0
<i>Pyrgo ringens</i>	0	30	0	0	0	0	0	0	0	100	0	0
<i>Pyrgo subsphaerica</i>	60	30	0	0	0	0	0	0	225	0	0	0
<i>Quinqueloculina intricata</i>	0	60	0	80	0	0	0	180	0	200	0	90
<i>Quinqueloculina lamarckiana</i>	2250	1290	5175	5080	19	441	8820	5160	225	4800	371	1395
<i>Quinqueloculina limbata</i>	0	0	0	0	0	0	0	420	0	450	0	45
<i>Quinqueloculina milletti</i>	0	0	0	0	0	0	0	0	0	200	0	0
<i>Quinqueloculina patagonica</i>	1920	1680	3975	6040	22	247	11520	4620	225	2100	237	495
<i>Quinqueloculina sp.</i>	0	120	0	400	0	0	0	60	0	500	0	45
<i>Quinqueloculina tubilocula</i>	0	0	0	0	0	0	0	0	0	100	0	0
<i>Quinqueloculina undulata</i>	0	0	0	40	0	0	0	0	0	50	0	0
<i>Reussella sp.</i>	0	0	25	0	0	0	0	0	45	300	0	0
<i>Rosalina sp.</i>	0	30	0	0	0	0	0	0	1575	0	0	0
<i>Siphogenerina raphana</i>	0	0	0	40	0	0	120	0	0	0	0	0
<i>Siphogenerina striatula</i>	0	0	0	0	0	0	0	0	0	50	0	0
<i>Siphonina bradyana</i>	0	0	0	160	0	0	0	0	0	200	0	45
<i>Spirulina vivipara</i>	0	0	0	0	0	0	0	0	0	50	0	0
<i>Spiroculina sp.</i>	0	0	0	0	0	0	0	0	0	200	0	90
<i>Spiroculina subimpresca</i>	0	0	0	40	0	0	0	0	0	50	0	0
<i>Textularia earlandii</i>	0	150	50	560	0	17	1380	60	45	350	25	0
<i>Textularia gramen</i>	300	270	1000	1400	0	13	1560	420	0	750	66	270
<i>Textularia sp.</i>	0	0	0	280	0	0	0	60	0	100	0	0
<i>Triloculina baldai</i>	0	930	0	3600	0	0	0	2520	1035	2100	0	585
<i>Uvigerina peregrina</i>	0	0	25	0	0	12	780	60	0	50	0	0
<i>Uvigerina striata</i>	0	0	0	0	0	0	0	120	0	0	0	0
<i>Wiesnerella sp.</i>	0	30	0	200	0	0	0	240	0	150	0	0
<i>Wiesnerella ujjei</i>	0	0	0	0	0	0	0	180	0	50	0	0

Source: authors

The groups shown by the cluster evident groups formed by the oldest (9 and 10), (6, 7, 8 and 11) (4, 5, and 12) most recent stations (1, 2, 3). The grouping is not clear; we still need further analysis (Figure 8).

Figure 8 – Cluster analysis among species of foraminifera and analysis of MDS in the intervals of core 3.



Source: authors

Table 6 shows the number of species varied between 7 and 60 and the number of species varied between 152 and 62040 individuals. Evenness showed a lower value (0.51) in the interval 3 and higher value (0.797951) in


the interval 9. Diversity was lower in the interval 5 (1.42) and highest in the interval 10 (2.853171992). The dominance showed the lowest value (0.11) in the interval 9 and the highest value in the interval 7 (0.34).

Table 6 – Number of species, specimens, evenness, diversity and dominance

Intervals	Species number	Specimens number	Evenness	Diversity	Dominance
1 (topo)	3	520	0.59	1.52	0.29
2	4	14790	0.62	2.21	0.23
3	20	200	0.51	1.52	0.33
4	41	54640	0.56	2.10	0.25
5	7	152	0.73	1.42	0.32
6	14	1702	0.63	1.68	0.25
7	17	62040	0.52	1.48	0.34
8	46	38820	0.61	2.35	0.19
9	13	5490	0.79	2.04	0.16
10	60	37050	0.69	2.85	0.12
11	15	1473	0.63	1.71	0.25
12 (base)	26	12240	0.60	1.97	0.26

Source: authors

Figure 9 – Diversity, dominance and evenness of Core 3



Source: authors

4 DISCUSSION

This study investigated the ecologically fragile region of an inner shelf coral reef through foraminifers' species in the last 25 years approximately. Closer to River Pium we observe greater oscillation due to intrannual variations of drought and rain. A little southern the environment was more diverse and stable with lower dominances of species. We observed intrannual fluctuations in three different microenvironments, but there is a declining trend in diversity and evenness, with greater dominance of few species nowadays. Through the study of benthic foraminifera population dynamics, we have evidenced a decrease in the diversity and evenness, and increase in dominance today when compared with values of 25 years ago in 3 distinct regions in the reef area.

In a study of a coral reef supporting carbonate shelf, Almeida¹⁰ has found three bio sedimentary units or functional groups¹¹: symbiotic-bearing foraminifers (*Amphistegina*, *Peneroplis* and *Archais*), other small taxa (*Miliolinella* and *Cornuspira*) and stress tolerant genera (*Bolivina*, *Elphidium* and *Ammonia*). The same author states that the reef degradation over time can be estimated by shifts in the FORAM index that are based on grouping foraminifera into the above-mentioned functional groups. Our study reveal that the absence of symbiotic-bearing foraminifers and the dominance of stress tolerant genera, which in according to Almeida¹² is due to the loss of favorable environmental conditions to support a healthy fauna even in the past. In according to Uthicke¹³ studying benthic Foraminifera as ecological indicators for water quality on the Great Barrier Reef revealed that low light and higher nutrient conditions is acceptable for the stress tolerant heterotrophic Rotaliida but several large foraminifera symbiotic-bearing ones were identified as indicators for offshore, clear water conditions. Our study revealed that the loss of environmental quality is evident by the decrease of biodiversity towards today.

10 ALMEIDA, C.M.; BARBOSA, C.F.; CORDEIRO, R.C.; SEOANE, J.C.S.; FERMINO, G.M.; SILVA, P.O.; TURCO, B.J. Palaeoecology of a 3-kyr biosedimentary record of a coral reef-supporting carbonate shelf. *Continental Shelf Research*, n. 70, p.168-176, 2013.

11 HALLOCK, P. Symbiont-bearing foraminifera: harbingers of global change, *Micropaleontology*, n.46, Suppl. 1, p.95-104. Google Scholar.

12 Op. cit.

13 UTHICKE, S.; NOBES, K. Benthic Foraminifera as ecological indicators for water quality on the Great Barrier Reef. *Estuarine, Coastal and Shelf Science*, v. 78, Issue 4, p. 763-773, 2008. ISSN 0272-7714, <http://dx.doi.org/10.1016/j.eccs.2008.02.014>.

REFERENCES

- ALMEIDA, C.M.; BARBOSA, C.F.; CORDEIRO, R.C.; SEOANE, J.C.S.; FERMINO, G.M.; SILVA, P.O.; TURCO, B.J. Palaeoecology of a 3-kyr biosedimentary record of a coral reef-supporting carbonate shelf. *Continental Shelf Research*, n. 70, p.168-176, 2013.
- EICHLER, P.P.B.; BILLUPS, K.; VELEZ, C.C. Investigating faunal and geochemical methods for tracing salinity in an Atlantic Coastal lagoon, Delaware, USA. *J. For. Res.*, v. 40, p. 14-33, 2010.
- EICHLER, P.P.B.; EICHLER, B.B.; MIRANDA, L.B. de; RODRIGUES, A.R. Foraminiferal assemblages in a subtropical, mixohaline, estuarine channel, Bertioga (São Paulo). *Braz. J. Res.*, n. 37, p. 45-58, 2007.
- EICHLER, P.P.B.; EICHLER, B.B.; SEN GUPTA, B.K.; RODRIGUES, A.R. Foraminifera as indicators of marine pollutant contamination on the inner continental shelf of southern Brazil. *Marine Pollution Bulletin*, v. 64, p. 22-30, 2012.
- EICHLER, P.P.B.; SEN GUPTA, B.K.; EICHLER, B.B.; BRAGA, E.S.; CAMPOS, E.J. Benthic foraminiferal assemblages of the South Brazil: Relationship to water masses and nutrient distributions. *Cont. Shelf. Res.*, n. 28, p. 1674-1686, 2008.
- HALLOCK, P. Symbiont-bearing foraminifera: harbingers of global change. *Micropaleontology*, n. 46, Suppl. 1, p. 95-104, 2000.
- PATCHINEELAM, S.R.; SMOAK, J.M. Sediment accumulation rates along the inner eastern Brazilian continental shelf. *Geo-Marine Letters*, n. 19, p. 196-201. 1999.
- SEN GUPTA, B.K. *Modern Foraminifera*. New York: Kluwer Academic Publishers, 2003.
- SEN GUPTA, B.K.; PLATON, E. Tracking past sedimentary records of oxygen depletion in coastal waters: use of the Ammonia-Elphidium Foraminiferal Index. *Journal of Coastal Research, Special Issue*, v.39, n. 8-12, p.1351-1355, 2006.
- UTHICKE, S.; NOBES, K. Benthic Foraminifera as ecological indicators for water quality on the Great Barrier Reef. *Estuarine, Coastal and Shelf Science*, v. 78, Issue 4, pages 763-773, ISSN 0272-7714. Disponível em: <<http://dx.doi.org/10.1016/j.ecss.2008.02.014>>.
- YANKO, V. Benthic foraminifera as bioindicators of stress environment: anthropogenic problems – foraminiferal solution. In: *The First International Conference Application of Micropaleontology in Environmental Sciences*. Tel Aviv Program & Abstracts, Herzlia: Anamet Ltd, 1997. p.117-119.
- YANKO, V.; AHMAD, M.; KAMINSKY, M. Morphological deformities of benthic foraminiferal tests in response to pollution by heavy metals: implications for pollution monitoring. *J. For. Res.*, v. 28, p. 177-200, 1998.
- YANKO, V.; ARNOLD, A.J.; PARKER, W.C. Effects of marine pollution on benthic Foraminifera In: SEN GUPTA, B.K. (Ed.). *Modern Foraminifera*. New York: Kluwer Acad. Publ., 1999. p.217-235.
- YANKO, V.; KRONFELD, J.; FLEXER, A. Response of benthic foraminifera to various pollution sources: implications for pollution monitoring. *J. For. Res.*, v. 24, p. 1-17, 1994.